

## Definitions

$$\bullet \cos A = \frac{\text{adjacent}}{\text{hypotenuse}} = \frac{x}{r}$$



$$\bullet \sin A = \frac{\text{opposite}}{\text{hypotenuse}} = \frac{y}{r}$$

$$\bullet \tan A = \frac{\text{opposite}}{\text{adjacent}} = \frac{y}{x}$$

$$\bullet \sec A = \frac{\text{hypotenuse}}{\text{adjacent}} = \frac{r}{x}$$


$$\bullet \csc A = \frac{\text{hypotenuse}}{\text{opposite}} = \frac{r}{y}$$

$$\bullet \cot A = \frac{\text{adjacent}}{\text{opposite}} = \frac{x}{y}$$



	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$90^\circ$
	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
tan	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	undefined

## Unit Circle



## Pythagorean Identities

- $\bullet \cos^2 A + \sin^2 A = 1$
- $\bullet 1 + \tan^2 A = \sec^2 A$
- $\bullet 1 + \cot^2 A = \csc^2 A$

## Reciprocal Identities

- $\bullet \sec A = \frac{1}{\cos A}$
- $\bullet \csc A = \frac{1}{\sin A}$
- $\bullet \cot A = \frac{1}{\tan A}$

## Ratio Identities

- $\bullet \tan A = \frac{\sin A}{\cos A}$
- $\bullet \cot A = \frac{\cos A}{\sin A}$

## Cofunction Identities

- $\bullet \cos(\frac{\pi}{2} - A) = \sin A$
- $\bullet \sin(\frac{\pi}{2} - A) = \cos A$
- $\bullet \tan(\frac{\pi}{2} - A) = \cot A$
- $\bullet \sec(\frac{\pi}{2} - A) = \csc A$
- $\bullet \csc(\frac{\pi}{2} - A) = \sec A$
- $\bullet \cot(\frac{\pi}{2} - A) = \tan A$

## Even/Odd Identities

- $\bullet \cos(-A) = \cos A$
- $\bullet \sin(-A) = -\sin A$
- $\bullet \tan(-A) = -\tan A$
- $\bullet \sec(-A) = \sec A$
- $\bullet \csc(-A) = -\csc A$
- $\bullet \cot(-A) = -\cot A$

## Sum and Difference Identities

- $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$
  - $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$
  - $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$
- 

## Double Angle Identities

- $\cos 2A = \cos^2 A - \sin^2 A$
  - $\sin 2A = 2 \cos A \sin A$
  - $\cos 2A = 2 \cos^2 A - 1$
  - $\cos 2A = 1 - 2 \sin^2 A$
  - $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$
- 

## Half Angle Identities

- $\cos \frac{A}{2} = \pm \sqrt{\frac{1 + \cos A}{2}}$
  - $\sin \frac{A}{2} = \pm \sqrt{\frac{1 - \cos A}{2}}$
  - $\tan \frac{A}{2} = \frac{1 - \cos A}{\sin A} = \frac{\sin A}{1 + \cos A}$
- 

## Power Reduction Identities

- $\cos^2 A = \frac{1 + \cos 2A}{2}$
  - $\sin^2 A = \frac{1 - \cos 2A}{2}$
  - $\tan^2 A = \frac{1 - \cos 2A}{1 + \cos 2A}$
- 

## Sum-to-Product Identities

- $\sin A + \sin B = 2 \sin \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right)$
  - $\sin A - \sin B = 2 \cos \left( \frac{A+B}{2} \right) \sin \left( \frac{A-B}{2} \right)$
  - $\cos A + \cos B = 2 \cos \left( \frac{A+B}{2} \right) \cos \left( \frac{A-B}{2} \right)$
  - $\cos A - \cos B = -2 \sin \left( \frac{A+B}{2} \right) \sin \left( \frac{A-B}{2} \right)$
- 

## Product-to-Sum Identities

- $\sin A \cos B = \frac{1}{2} [\sin(A+B) + \sin(A-B)]$
  - $\cos A \cos B = \frac{1}{2} [\cos(A-B) + \cos(A+B)]$
  - $\sin A \sin B = \frac{1}{2} [\cos(A-B) - \cos(A+B)]$
- 

## Sums of Sines and Cosines

- $A \cos x + B \sin x = \sqrt{A^2 + B^2} \sin(x + \phi)$  where  $\cos \phi = \frac{B}{\sqrt{A^2 + B^2}}$  and  $\sin \phi = \frac{A}{\sqrt{A^2 + B^2}}$
  - $A \cos x + B \sin x = \sqrt{A^2 + B^2} \cos(x - \phi)$  where  $\cos \phi = \frac{A}{\sqrt{A^2 + B^2}}$  and  $\sin \phi = \frac{B}{\sqrt{A^2 + B^2}}$
- 

## Circular Sections

- Arc length:  $s = r\theta$
  - Area:  $A = \frac{1}{2}r^2\theta$
  - Angular velocity:  $\omega = \frac{v}{r}$
- 

## Graphing

For  $y = A \cos(Bx + C) + D$  and  $y = \sin(Bx + C) + D$ ,

- Amplitude =  $|A|$
- Frequency =  $B$
- Vertical Shift =  $D$
- Period =  $\frac{2\pi}{B}$
- Phase Shift =  $-\frac{C}{B}$

The graph of  $y = \tan(Bx + C)$  has asymptotes at the locations where  $\cos(Bx + C) = 0$ .

---

## Range of Inverse Functions

- $y = \arccos x, \quad 0 \leq y \leq \pi$
  - $y = \arcsin x, \quad -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$
  - $y = \arctan x, \quad -\frac{\pi}{2} < y < \frac{\pi}{2}$
  - $y = \operatorname{arcsec} x, \quad 0 \leq y < \frac{\pi}{2} \text{ and } \frac{\pi}{2} < y \leq \pi$
  - $y = \operatorname{arccsc} x, \quad -\frac{\pi}{2} \leq y < 0 \text{ and } 0 < y \leq \frac{\pi}{2}$
  - $y = \operatorname{arccot} x, \quad -\frac{\pi}{2} < y < 0 \text{ and } 0 < y < \frac{\pi}{2}$
- 

## Laws of Sines and Cosines

- $c^2 = a^2 + b^2 - 2ab \cos C$
  - $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
-