

Factoring

- $a^2 - b^2 = (a - b)(a + b)$
 - $a^2 + b^2$ is prime
 - $a^2 + 2ab + b^2 = (a + b)^2$
 - $a^2 - 2ab + b^2 = (a - b)^2$
 - $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$
 - $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$
-

Analytic Geometry

- slope: $m = \frac{y_2 - y_1}{x_2 - x_1}$
 - equation of a line: $y - y_1 = m(x - x_1)$
 - distance: $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
-

Exponent Rules

- $a^{x+y} = a^x a^y$
 - $(ab)^x = a^x b^x$
 - $(a^x)^y = a^{xy}$
 - $a^0 = 1$ if $a \neq 0$
 - $a^{-x} = \frac{1}{a^x}$ if $a \neq 0$
 - $a^{x-y} = \frac{a^x}{a^y}$ if $a \neq 0$
-

Logarithm Rules

- $\log_b x = y \iff x = b^y$
 - $b^{\log_b x} = x$
 - $\log_b b^x = x$
 - $\log_b 1 = 0$
 - $\log_b b = 1$
 - $\log_b xy = \log_b x + \log_b y$
 - $\log_b \frac{x}{y} = \log_b x - \log_b y$
 - $\log_b x^y = y \log_b x$
 - $\log_b x = \frac{\log_a x}{\log_a b}$
-

Arithmetic Series

- $a_k = a + (k - 1)d$
 - $S_n = \sum_{k=1}^n [a + (k - 1)d] = \frac{n}{2} [2a + (n - 1)d]$
 - $S_n = \sum_{k=1}^n [a + (k - 1)d] = n \left(\frac{a + a_n}{2} \right)$
-

Geometric Series

- $a_n = ar^{n-1}$
 - $S_n = \sum_{k=0}^{n-1} ar^k = a \left[\frac{1 - r^n}{1 - r} \right]$ if $r \neq 1$
 - $S = \sum_{k=0}^{\infty} ar^k = \frac{a}{1 - r}$ if $|r| < 1$
-

Trigonometry

- $\cos A = \frac{\text{adjacent}}{\text{hypotenuse}}$ • $\sin A = \frac{\text{opposite}}{\text{hypotenuse}}$
- $\tan A = \frac{\text{opposite}}{\text{adjacent}}$

	0°	30°	45°	60°	90°
	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
tan	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	undefined

Pythagorean Identities

- $\cos^2 A + \sin^2 A = 1$
 - $1 + \tan^2 A = \sec^2 A$
 - $1 + \cot^2 A = \csc^2 A$
-

Ratio Identities

- $\tan A = \frac{\sin A}{\cos A}$ • $\cot A = \frac{\cos A}{\sin A}$
-

Reciprocal Identities

$$\bullet \sec A = \frac{1}{\cos A} \quad \bullet \csc A = \frac{1}{\sin A} \quad \bullet \cot A = \frac{1}{\tan A}$$

Sum and Difference Identities

$$\begin{aligned} \bullet \cos(A \pm B) &= \cos A \cos B \mp \sin A \sin B \\ \bullet \sin(A \pm B) &= \sin A \cos B \pm \cos A \sin B \\ \bullet \tan(A \pm B) &= \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \end{aligned}$$

Double Angle Identities

$$\begin{aligned} \bullet \cos 2A &= \cos^2 A - \sin^2 A \\ \bullet \cos 2A &= 2 \cos^2 A - 1 \\ \bullet \cos 2A &= 1 - 2 \sin^2 A \\ \bullet \sin 2A &= 2 \cos A \sin A \\ \bullet \tan 2A &= \frac{2 \tan A}{1 - \tan^2 A} \end{aligned}$$

Half Angle Identities

$$\begin{aligned} \bullet \cos \frac{A}{2} &= \pm \sqrt{\frac{1 + \cos A}{2}} \\ \bullet \sin \frac{A}{2} &= \pm \sqrt{\frac{1 - \cos A}{2}} \\ \bullet \tan \frac{A}{2} &= \frac{1 - \cos A}{\sin A} = \frac{\sin A}{1 + \cos A} \end{aligned}$$

Triple Angle Identities

$$\begin{aligned} \bullet \cos 3A &= 4 \cos^3 A - 3 \cos A \\ \bullet \sin 3A &= 3 \sin A - 4 \sin^3 A \end{aligned}$$

Power Reduction Identities

$$\begin{aligned} \bullet \cos^2 A &= \frac{1 + \cos 2A}{2} \\ \bullet \sin^2 A &= \frac{1 - \cos 2A}{2} \\ \bullet \tan^2 A &= \frac{1 - \cos 2A}{1 + \cos 2A} \\ \bullet \cos^3 A &= \frac{3 \cos A + \cos 3A}{4} \\ \bullet \sin^3 A &= \frac{3 \sin A - \sin 3A}{4} \end{aligned}$$

Sum-to-Product Identities

$$\begin{aligned} \bullet \sin A + \sin B &= 2 \sin \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) \\ \bullet \sin A - \sin B &= 2 \cos \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right) \\ \bullet \cos A + \cos B &= 2 \cos \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) \\ \bullet \cos A - \cos B &= -2 \sin \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right) \end{aligned}$$

Product-to-Sum Identities

$$\begin{aligned} \bullet \sin A \cos B &= \frac{1}{2} [\sin(A+B) + \sin(A-B)] \\ \bullet \cos A \cos B &= \frac{1}{2} [\cos(A+B) + \cos(A-B)] \\ \bullet \sin A \sin B &= \frac{1}{2} [\cos(A-B) - \cos(A+B)] \end{aligned}$$

Sums of Sines and Cosines

$$\begin{aligned} \bullet A \cos x + B \sin x &= \sqrt{A^2 + B^2} \sin(x + \phi) \text{ where} \\ &\cos \phi = \frac{B}{\sqrt{A^2 + B^2}} \text{ and } \sin \phi = \frac{A}{\sqrt{A^2 + B^2}} \\ \bullet A \cos x + B \sin x &= \sqrt{A^2 + B^2} \cos(x - \phi) \text{ where} \\ &\cos \phi = \frac{A}{\sqrt{A^2 + B^2}} \text{ and } \sin \phi = \frac{B}{\sqrt{A^2 + B^2}} \end{aligned}$$

Laws of Sines and Cosines

$$\begin{aligned} \bullet c^2 &= a^2 + b^2 - 2ab \cos C \\ \bullet \frac{a}{\sin A} &= \frac{b}{\sin B} = \frac{c}{\sin C} \end{aligned}$$

Area of a Triangle

For a triangle with sides a , b , c and angles $\angle A$, $\angle B$, and $\angle C$,

$$\begin{aligned} \bullet \text{Area} &= \frac{\sqrt{s(s-a)(s-b)(s-c)}}{2} \text{ where} \\ &s = \frac{a+b+c}{2} \\ \bullet \text{Area} &= \frac{1}{2} ab \sin C \\ \bullet \text{Area} &= \frac{c^2 \sin A \sin B}{2 \sin C} \end{aligned}$$

Circular Section

$$\begin{aligned} \bullet \text{Arc length: } &s = r\theta \\ \bullet \text{Area: } &A = \frac{1}{2} r^2 \theta \end{aligned}$$
